

IOWA STATE UNIVERSITY

Digital Repository

Beef Research Report, 2001

Animal Science Research Reports

2002

Genetic Relationship between Live Animal Ultrasound Measures and Scrotal Circumference in Yearling Angus Bulls

Doyle E. Wilson
Iowa State University

Gene H. Rouse
Iowa State University

Craig L. Hays
Iowa State University

Follow this and additional works at: http://lib.dr.iastate.edu/beefreports_2001



Part of the [Animal Sciences Commons](#)

Extension Number: ASL R1737

Recommended Citation

Wilson, Doyle E.; Rouse, Gene H.; and Hays, Craig L., "Genetic Relationship between Live Animal Ultrasound Measures and Scrotal Circumference in Yearling Angus Bulls" (2002). *Beef Research Report, 2001*. 3.
http://lib.dr.iastate.edu/beefreports_2001/3

This report is brought to you for free and open access by the Animal Science Research Reports at Iowa State University Digital Repository. It has been accepted for inclusion in Beef Research Report, 2001 by an authorized administrator of Iowa State University Digital Repository. For more information, please contact digirep@iastate.edu.

Genetic Relationship between Live Animal Ultrasound Measures and Scrotal Circumference in Yearling Angus Bulls

Abstract

The purpose of this study was to determine the relationship between 365-d yearling scrotal circumference, scanning weight, and component ultrasound measured compositional traits. The genetic correlations between scrotal circumference measured at 365-d of age and other live animal measures are low. Of these five genetic relationships, the highest correlation is with % intramuscular fat (IMF) at .13. However, this is a low genetic correlation. The positive sign of the scrotal circumference-% IMF genetic correlation is encouraging for breeders wanting to increase scrotal circumference and % IMF at the same time. Scrotal circumference in yearling Angus bulls is not strongly linked genetically with weight and compositional traits. Selection programs for either increased retail product (as measured by weight and ribeye area) or quality (as measured by % IMF) will not result in an antagonistic effect on scrotal circumference.

Keywords

ASL R1737

Disciplines

Animal Sciences

Genetic Relationship between Live Animal Ultrasound Measures and Scrotal Circumference in Yearling Angus Bulls

A.S. Leaflet R1737

Doyle E. Wilson, professor of animal science
Gene H. Rouse, professor of animal science
Craig L. Hays, mgr., centralized
ultrasound-processing laboratory

Summary

The purpose of this study was to determine the relationship between 365-d yearling scrotal circumference, scanning weight, and component ultrasound measured compositional traits. The genetic correlations between scrotal circumference measured at 365-d of age and other live animal measures are low. Of these five genetic relationships, the highest correlation is with % intramuscular fat (IMF) at .13. However, this is a low genetic correlation. The positive sign of the scrotal circumference-% IMF genetic correlation is encouraging for breeders wanting to increase scrotal circumference and % IMF at the same time. Scrotal circumference in yearling Angus bulls is not strongly linked genetically with weight and compositional traits. Selection programs for either increased retail product (as measured by weight and ribeye area) or quality (as measured by % IMF) will not result in an antagonistic effect on scrotal circumference.

Introduction

Real-time ultrasound has opened the door to opportunities for studying growth and development in beef cattle as never before. This tool is being used to annually measure thousands of young seed stock animals directly to determine genetic merit for compositional traits. The indications are that yearling bulls can be scanned for traits of external fat, ribeye area, and percentage intramuscular fat

(% IMF) to accurately predict these same traits in steer progeny destined for harvest. One of the issues that is unresolved is the relationship between what is happening to lean and fat deposition when the young bull begins to mature sexually. Particularly questioned is whether the process of achieving sexual maturity has any effect on the deposition or levels of % IMF that could potentially bias carcass expected progeny differences (EPD) based ultrasound measures. The purpose of this study was to determine the genetic relationship between 365-d yearling scrotal circumference, scanning weight, and component ultrasound measured compositional traits.

Materials and Methods

Twice annually, Iowa State University (ISU) conducts a genetic evaluation of scrotal circumference in yearling bulls for the American Angus Association. ISU is also conducting a genetic evaluation for ultrasound measured traits in yearling Angus bulls and developing Angus heifers. For the Spring-2000 evaluation, there were 128,518 scrotal records and 29,938 yearling bull ultrasound records. There were 11,852 yearling bulls having both scrotal measures and ultrasound measures, and these animals were used to conduct this study. There were 1,191 sires represented by these yearling bulls. Both scrotal and ultrasound measures were adjusted to a 365-d age end point and for age of dam effects. A summary of the adjusted records is presented in Table 1.

A multiple-trait sire-maternal grandsire model was used to estimate the genetic variance-covariance parameters using restricted maximum likelihood procedures. Contemporary group effects fit in the same model included herd, weaning contemporary group, and scanning contemporary group.

Table 1. Adjusted yearling Angus bull live scan and scrotal circumference measures (11,852 head)

| Trait | Mean | Std. Dev. | Minimum | Maximum |
|----------------------------------|-------|-----------|---------|---------|
| Scan weight, lbs | 1075 | 110 | 587 | 1553 |
| Gain, lbs/d | 2.95 | .55 | .31 | 5.67 |
| % IMF, % | 3.42 | .82 | 1.07 | 8.49 |
| Ribeye area, sq. in. | 12.06 | 1.38 | 6.00 | 17.7 |
| 12-13 th rib fat, in. | .25 | .10 | .03 | .93 |
| Rump fat, in. | .30 | .10 | .02 | .87 |
| Scrotal cir., cm | 36.2 | 2.6 | 23.1 | 49.0 |

Results and Discussion

The heritability, genetic correlation and phenotypic correlation estimates are presented in Table 2. The Heritability estimates are consistent with estimates

previously derived from the independent data sets (ultrasound and scrotal).

The genetic correlations between scrotal circumference measured at 365-d of age and other live animal measures are

2001 Beef Research Report — Iowa State University

low. Of the five genetic relationships, the highest is with % IMF at .13. However, this is a low genetic correlation. The positive sign of the scrotal circumference-% IMF genetic correlation is encouraging for breeders wanting to increase scrotal circumference and % IMF at the same time. The largest correlation is a phenotypic correlation with scanning weight at .27. This correlation suggests larger bulls are going to have larger testicles at a year of age. The other phenotypic correlations are small, but positive.

The next step in more fully understanding the sexual maturity and growth and development patterns in yearling

Angus bulls will be to study within-animal growth curves for each of these two traits. This research is ongoing at ISU as part of the beef cattle breeding project. Serial measurements of these traits from weaning to harvest are being collected annually on intact males starting with the 1998 calf crop. The first results from an analysis of three years of data will be available mid-summer 2001.

It should be noted that the h^2 level for scrotal circumference is in the moderate range (.34). With a h^2 of this magnitude, selection can be used to increase (or decrease) circumference.

Table 2. Heritability, genetic correlation and phenotypic correlation estimates for yearling Angus bull live scan and scrotal circumference measures (365-d age end point).

| Trait | SWT | % IMF | REA | FAT | SCR | RFAT |
|-----------------------------------|------------------------|-------------------|-----------|-----------|-----------|-----------|
| Scan weight (SWT) | .57 ^a (.06) | -.16 ^b | .30 | .11 | .05 | .12 |
| % IMF | .00 ^c | .32 (.05) | -.07 | .24 | .13 | .16 |
| Ribeye area (REA) | .40 | -.07 | .34 (.05) | .20 | .05 | .10 |
| 12-13 th rib fat (FAT) | .30 | .19 | .19 | .37 (.06) | .06 | .66 |
| Scrotal (SCR) | .27 | .03 | .12 | .12 | .34 (.05) | .06 |
| Rump fat (RFAT) | .24 | .16 | .13 | .60 | .11 | .36 (.06) |

^aHeritability with standard error in parenthesis.

^bGenetic correlations on the upper off-diagonals.

^cPhenotypic correlations on the lower off-diagonals.

Implications

Scrotal circumference in yearling Angus bulls is not strongly linked genetically with weight and compositional traits. Selection programs for either increased retail product (as measured by weight and ribeye area) or quality (as measured by %

IMF) will not result in an antagonistic effect on scrotal circumference.

Acknowledgments

The authors express their appreciation to the American Angus Association for the financial support of this research and the use of their data to conduct the analysis.